## South Fork Payette River Subbasin Assessment



Final



Department of Environmental Quality

July 2005

### **Subbasin Assessment of the South Fork Payette River**

**July 2005** 

Prepared by:
Craig Shepard
Boise Regional Office
Department of Environmental Quality
1445 North Orchard
Boise, Idaho 83706

### **Acknowledgments**

This subbasin assessment would not have been possible without the efforts and resources of the following individuals and agencies:

Craig Shepard, Charlie Bidondo, Bryan Horsburgh and Angie Petersen of the Idaho Department of Environmental Quality, Boise Regional Office, provided administrative resources, water body assessments, water quality data and background literature pertinent to the South Fork Payette River.

The Boise National Forest and the Sawtooth National Forest provided staff time and data. T. J. Clifford provided technical references, coordinated Boise National Forest resources and assisted with development of an assessment strategy. John Thornton and Carey Crist provided spatial data from the Boise National Forest and the Southwest Idaho Ecogroup. Michael Kellett provided biological and water quality data. Susy Osgood supplied historical and cultural references. Valdon Hancock provided technical references from the Sawtooth National Forest.

Don Zaroban drafted the original document and the Boise Regional Office including Bryan Horsburgh, Mike Ingham, Marti Bridges and Craig Shepard provided review and additional information for the final document. Mary Grandjean supplied historical and cultural information. Sean Coyle prepared maps of the spatial data. Dennis Meier and Charles Huntington provided technical editing assistance.

Water quality and stream discharge data were downloaded from the United States Geological Survey (USGS) Web site. Climate data were downloaded from the Western Regional Climate Center World Wide Web site.

The cover photograph, taken by Cynthia Grafe on August 31, 1998, shows the South Fork Payette River near Lowman, Idaho.

## **Table of Contents**

Acknowledgments	i
Table of Contents	iii
List of Tables	v
List of Figures	viii
Abbreviations, Acronyms, and Symbols	xi
Executive Summary	xiii
Subbasin at a Glance	xiii
Key Findings	xiv
Subbasin Assessment – Watershed Characterization	1
1.1 Introduction	
Background	1
Idaho's Role	2
1.2 Physical and Biological Characteristics	
Climate	
Subbasin Characteristics	8
Subwatershed and Stream Characteristics	16
1.3 Cultural Characteristics	19
Land Use	19
Land Ownership, Cultural Features, and Population	
History and Economics	23
2. Subbasin Assessment – Water Quality Concerns and Status	27
2.1 Water Quality Limited Segments Identified in the Subbasin	
2.2 Applicable Water Quality Standards	
Beneficial Uses	30
Existing Uses	30
Designated Uses	30
Presumed Uses	31
Pollutant Relationships to Uses – Target Identification	32
2.3 Summary and Analysis of Existing Water Quality Data	
South Fork Payette River	34
6.4.2. River Index Scoring	42
Other Water Bodies	
Chapman Creek	49

Smokey Creek	57
Horn Creek	
Wash Creek	73
Conclusions	80
2.4 Data Gaps	
3. Subbasin Assessment – Pollutant Source Inventor	
3.1 Sources of Pollutants of Concern	
Point Sources	
Nonpoint Sources	83
Pollutant Transport	84
Estimation and Prioritization of Sedimen	nt Sources84
Land Type Erosion Hazard	84
Road Density	
Percent Surface Fines	86
Fires and Landslides	89
Source Summary	92
4. Pollution Control Efforts Subbasin Assessme and Status	95
Boise National Forest	
Idaho Forest Practices Act	
Idaho Transportation Department	
idano Transportation Department	
References Cited	99
Glossary	103
Appendices	113
Appendix A. Unit Conversion Chart	
Appendix B. State and Site-Specific Standar	
Appendix C. Data Sources	121
Appendix D. Excerpts from the Boise Nation	
Appendix E. Statistical Analysis for Normali	
Appendix F. Distribution List	
Annendix G. Public Comments	131

### **List of Tables**

Table A. Summary of assessment outcomes	χv
Table 1. Summary of period of record, annual maximum and minimum air	
temperatures and total annual precipitation for Garden Valley, Lowman	
and Deadwood Dam climate stations	5
Table 2. Common name, scientific name and status of origin of fish native to	,
or introduced (Zaroban et al. 1999) in the South Fork Payette River	
subbasin	16
Table 3. Surface area (acres) of the 35 HUC6 watersheds in the South Fork	
Payette River subbasin	
Table 4. Estimates of the minimum stream elevation (feet), maximum stream	
elevation (feet), segment length (miles) and drop in elevation (feet per	
mile) for the mainstem stream	18
Table 5. Water Quality Limited Segments in the South Fork Payette River	
subbasin	28
Table 6. South Fork Payette River subbasin designated beneficial uses	30
Table 7. South Fork Payette River subbasin existing beneficial uses	31
Table 8. USGS surface water quality and quantity monitoring locations in the	<b>)</b>
South Fork Payette River Subbasin	34
Table 9. Suspended Sediment Concentration Results for South Fork Payette	<b>)</b>
at Lowman 1991, 1992, 1994, 1995, 1997, 1998 and 2001	<b>37</b>
Table 10. Measured and Estimated Discharge, Suspended Sediment Loads,	
Suspended Sediment Concentration, and Error Bias for South Fork	
Payette at Lowman	39
Table 11. Normalized Discharge and Estimated Average Suspended Sedime	nt
Concentration and Loads for South Fork Payette at Lowman	39
Table 12. Application of Bias to Normalized Discharge and Estimated Average	ge
Suspended Sediment Concentration and Loads for South Fork Payette a	ıt
Lowman	_
Table 13. Values of diatom association metrics for Idaho rivers in 1998	42
Table 14. Macroinvertebrates (RMI)	
Table 14a. RMI, RDI, RFI, and RPI Scoring and Rating Categories	
Table 15. USGS Monitoring South Fork Payette River @ Lowman <sup>1</sup>	
Table 16. Macroinvertebrate metrics and invertebrate river index (IRI) <sup>1</sup>	44
Table 17. Status of beneficial uses for the South Fork Payette River	45
Table 18. BURP results for wadeable streams in the South Fork Payette Rive	r
	47
Table 18 (Cont.). BURP results for wadeable streams in the South Fork Payer	tte
River basin	
Table 19. Summary of DEQ reconnaissance fish data collected in 2001	49
Table 20. Stream Macroinvertebrate Metrics and Results for Final SMI Score	
	50
Table 21. Stream Habitat Metrics and Results for Final SHI Score for Chapma	
Creek	
Table 22 Final Condition Rating for Changan Creek	51

Table 23. 1997 Percent Fines ≤ 6mm. BURP Site 1997SBOIC029, Chapman	
Creek	56
Table 24. 2004 Percent Fines ≤ 6mm. BURP Site 1997SBOIC029, Chapman	
Creek	
Table 25. Status of beneficial uses in the South Fork Payette River basin	57
Table 26. Stream Macroinvertebrate Metrics and Results for Final SMI Score	
for Smokey Creek Upper	
Table 27. Stream Habitat Metrics and Results for Final SHI Score for Smokey	,
Upper Creek	58
Table 28. Stream Macroinvertebrate Metrics and Results for Final SMI Score	
for Lower Smokey Creek	58
Table 29. Stream Habitat Metrics and Results for Final SHI Score for Lower	
Smokey Creek	58
Table 30. Final Condition Rating for Smokey Creek	
Table 31. 1997 Percent Fines ≤ 6mm. BURP Site 1997SWIROB41, Upper	-
Smokey Creek.	66
Table 32. 1997 Percent Fines ≤ 6mm. BURP Site 1997SWIROB40, Lower	
Smokey Creek.	66
Table 33. 2004 Percent Fines ≤ 6mm. BURP Site 1997SWIROB40, Lower	
Smokey Creek.	67
Table 34. Status of beneficial uses in Smokey Creek	
Table 35. Stream Macroinvertebrate Metrics and Results for Final SMI Score	O1
	68
Table 36. Stream Habitat Metrics and Results for Final SHI Score for Horn	UU
Creek	60
Table 37. Final Condition Rating for Horn Creek	
Table 38. 1997 Percent Fines ≤ 6mm. BURP Site 1997SWIROB391, Horn Cree	
	r. 72
Table 39. 2004 Percent Fines ≤ 6mm. BURP Site 1997SWIROB39, Horn Creek	
Table 40 Otation of bour field was a in Hams One als	
Table 40. Status of beneficial uses in Horn Creek	12
Table 41. Stream Macroinvertebrate Metrics and Results for Final SMI Score	
for Upper Wash Creek.	13
Table 42. Stream Habitat Metrics and Results for Final SHI Score for Upper	
Wash Creek	74
Table 43. Stream Macroinvertebrate Metrics and Results for Final SMI Score	
for Lower Wash Creek	74
Table 44. Stream Habitat Metrics and Results for Final SHI Score for Lower	
Wash Creek	
Table 45. Final Condition Rating for Wash Creek	
Table 46. 1997 Percent Fines ≤ 6mm. BURP Site 1997SWIROB41, Wash Creel	
Upper	<b>78</b>
Table 47. 1997 Percent Fines ≤ 6mm. BURP Site 1997SWIROB40, Wash Creel	<
Lower	
Table 48. 2004 Percent Fines ≤ 6mm. BURP Site 1997SWIROB40, Wash Creel	<
Lower.	79

Table 49. Status of beneficial uses in Wash Creek	79
Table 50. Summary of the South Fork Payette River Subbasin assessment	80
Table 51. Data gaps identified during development of the South Fork Payette	;
River Subbasin Assessment	81
Table 52. Summary of erosion hazard, gradient, roads, fine-grained sediment	t
and overall geomorphic risk score for HUC6 watersheds	87
Table 52 (Cont.). Summary of erosion hazard, gradient, roads, fine-grained	
sediment and overall geomorphic risk score for HUC6 watersheds	88
Table 53. Summary of erosion hazard, gradient, roads, fine-grained sediment	t
and overall geomorphic risk score for Impaired Water Bodies	88

# **List of Figures**

Figure A. South Fork Payette River Subbasin at a Glancexiv Figure 1. South Fork Payette River Subbasin4
Figure 2. Summary of Mean Monthly Maximum and Minimum Air Temperature at Garden Valley, Idaho6
Figure 3. Summary of Mean Monthly Maximum and Minimum Air Temperature at Lowman, Idaho6
Figure 4. Summary of Mean Monthly Maximum and Minimum Air Temperatures at Deadwood Dam, Idaho.
Figure 5. Summary of Average Monthly Total Precipitation at Garden Valley,  Lowman and Deadwood Dam, Idaho7
Figure 6. Average Monthly Discharge (cubic feet per second [cfs]) of the South Fork Payette River at Lowman (736 observations, June 1941 through September 2002) and Deadwood River below Dam (912 observations,
October 1926 through September 2002)8
Figure 7. Sixth Field HUCs in the South Fork Payette River Subbasin9
Figure 8. Geology of the South Fork Payette Subbasin11
Figure 9. Slope Map of the South Fork Payette River Subbasin12
Figure 10. Land Cover in the South Fork Payette River Subbasin14
Figure 11. Land Types in the South Fork Payette River Subbasin
Figure 12. Land Use of the South Fork Payette River Subbasin20
Figure 13. Land Ownership in the South Fork Payette River Subbasin22
Figure 14. Road Density within the South Fork Payette River Subbasin25
Figure 15. Water Quality Limited Segments within the South Fork Payette River
Subbasin
Figure 16. Annual Average Discharge at Lowman, Idaho since 1942 as
Compared to Period of Record Average Discharge of 861 cfs35
Figure 17. Mean Daily Discharge SF Payette River at Lowman, 1941 through
200435
Figure 18. Measured Sediment Concentrations, 1991 through 200136
Figure 19. South Fork Payette River at Lowman, Natural Log Suspended
Sediment Load as a Function of Discharge37
Figure 20. Sediment Concentrations, SF Payette River at Lowman Normalized
Discharge + or – 24%40
Figure 21. Chapman Creek 199752
Figure 22. Chapman Creek 199752
Figure 23. Chapman Creek 200453
Figure 24. Chapman Creek 200453
Figure 25. Chapman Creek Watershed. 10 Meter LANDSAT Imagery54
Figure 26. Chapman Creek Watershed. Roads55
Figure 27. Smokey Creek. 1997-98 LANDSAT Imagery61
Figure 28. Smokey Creek Lower 200462
Figure 29. Smokey Creek 200462
Figure 30. Smokey Creek 200463
Figure 31. Smokey Creek 2004

Figure 32. Smokey Creek (Upper) 2004	. 64
Figure 33. Road System in the Smokey Creek Watershed	65
Figure 34. Horn Creek 2004	.70
Figure 35. Horn Creek 2004	.71
Figure 36. Horn Creek 2004	.71
Figure 37. Upper Wash Creek 1997	.76
Figure 38. Upper Wash Creek 1997	.76
Figure 39. Upper Wash Creek 2004	. 77
Figure 40. Lower Wash Creek 1997	. 77
Figure 41. Lower Wash Creek 2004	. 78
Figure 42. Jack Creek blowout – Middle Fork Salmon River, September 17,	
2002	. 90
Figure 43. Orelano Creek blowout – Middle Fork Salmon River, September	
2002	. 90
Figure 44. Haystack Rapid – Middle Fork Salmon River, September 19, 2002	
(above)	. 91
Figure 45. Haystack Rapid – Middle Fork Salmon River, 2003 (below) after	
Bernard Creek blowout	. 91
Figure 46. Fifty Year Fire Occurrence in the South Fork Payette River	
Subbasin	. 93
Figure47. Locations of Significant 1995 Landslides in the South Fork Payette	•
River Subbasin	. 94
Figure 48. Highway 21 road cut	. 96
Figure 49. Highway 21 road cut	. 97
Figure 50. Highway 21 road cut	. 97
Figure 51. Highway 21 road cut	. 98

# Abbreviations, Acronyms, and Symbols

§303(d)	Refers to section 303	SBA	subbasin assessment
	subsection (d) of the Clean Water Act, or a list of	SCR	secondary contact recreation
	impaired water bodies required by this section		total maximum daily load
μ	Micro, one one-thousandth	TSS	total suspended solids
§	Section (usually a section of	U. S.	United States
federal or state rules or statutes)		U.S.C.	United States Code
AWS	agricultural water supply	USFS	United States Forest Service
C	Celsius	USGS	United States Geological Survey
cfs	cubic feet per second	WBID	water body identification number
cm	centimeters	WQLS	water quality limited segment
CWA	Clean Water Act	WQS	water quality standard
DWS	domestic water supply		
F	Fahrenheit		
HUC6	USGS sixth field (12-digit hydrologic unit code)		
IDAPA	Abbreviation for Idaho Administrative Procedures Act, refers to citations of Idaho administrative rules		
IDWR	Idaho Department of Water Resources		
NPDES	National Pollutant Discharge Elimination System		
NTU	nephelometric turbidity unit		
PCR	primary contact recreation		

### **Executive Summary**

The federal Clean Water Act (CWA) requires that states and tribes restore and maintain the chemical, physical, and biological integrity of the nation's waters. States and tribes, pursuant to Section §303 of the CWA, are to adopt water quality standards necessary to protect fish, shellfish, and wildlife while providing for recreation in and on the nation's waters whenever possible. Section 303(d) of the CWA establishes requirements for states and tribes to identify and prioritize water bodies that are water quality limited (i.e., water bodies that do not meet water quality standards). States and tribes must periodically publish a priority list (a"§303(d) list") of impaired waters. Currently this list must be published every two years. For waters identified on this list, states and tribes must develop a total maximum daily load (TMDL) for the pollutants, set at a level to achieve water quality standards.

This document addresses the water bodies in the South Fork Payette River Subbasin that have been placed on Idaho's current §303(d) list.


This subbasin assessment (SBA) has been developed to comply with Idaho's TMDL schedule. The assessment describes the physical, biological, and cultural setting; water quality status; pollutant sources; and recent pollution control actions in the South Fork Payette River Subbasin located in southwest Idaho.

The starting point for this assessment was Idaho's current §303(d) list of water quality limited water bodies. The SBA examines the current status of §303(d) listed waters and identifies potential sources of pollutants throughout the subbasin.

#### Subbasin at a Glance

From the Sawtooth Wilderness Area boundary to the confluence of the Middle Fork Payette River, the South Fork Payette River is listed on the 1998 §303(d) list for fine-grained sediment (Figure A). This listing was based on the 1990 Boise National Forest Management Plan (Boise National Forest 1990), which stated that a goal for the forest was to manage the watershed to reduce sedimentation in the South Fork Payette River. In the plan, beneficial uses, as identified in the Idaho water quality standards (WQS), were not reported as impaired.

The South Fork Payette River Subbasin contains bull trout, a species listed as threatened under the Endangered Species Act. The South Fork Payette River and Deadwood River are key bull trout watersheds in Idaho. Fine-grained sediment is one of the factors identified as limiting bull trout habitat in the subbasin.



#### **South Fork Payette River Basin**

5<sup>th</sup> Field HUC#: 17050120

Water Body: SF Payette River

Boundary: Wilderness Boundary to

Payette River

Ecoregion: Northern Mountains

Pollutants: Sediment

Beneficial Uses: Salmonid Spawning, Cold

Water Aquatic Life, Primary Contact Recreation

Pollution Sources: Non-Point Sources

Size: 520,318 acres

Figure A. South Fork Payette River Subbasin at a Glance.

#### **Key Findings**

The South Fork Payette River near Lowman, Idaho, has been monitored for water column sediment (suspended sediment concentration), bedload, and discharge. Extensive flow records also exist near Garden Valley, Idaho. Limited biological data are available for the South Fork Payette River. Most biological monitoring has been conducted in the tributaries.

The suspended sediment concentration data for the South Fork Payette River show that during years of normal flow, when mass wasting events are less prevalent, the water column sediment levels are below the 14-day duration suspended sediment target of 80 mg/L. The data also show that during years of high flow, erosion can be exacerbated and the sediment target exceeded. It is DEQ's belief that any excursions above the target at high flows are within the norms of natural variability and are not impairing beneficial uses.

To determine the expected suspended sediment concentration in the river in a typical flow year, a regression analysis was performed—between paired suspended sediment and flow data. The analysis determined that, at an average annual flow of 861 cfs, the suspended sediment concentration in the river would be 8.0 mg/L, well below the 14-day duration suspended sediment target of 80 mg/L. As a result of these analyses, DEQ does not recommend developing a sediment TMDL for the South Fork Payette River and recommends de-listing sediment from the §303(d) list.

While some anthropogenic sources of sediment (primarily from forest roads) exist within the basin and exceed the Boise National Forest desired future conditions during high flow events, DEQ believes this sediment is not currently impairing beneficial uses. However, road

management activities within the Boise and Sawtooth National Forests should be prioritized and maximized to reduce erosion. This strategy is outlined in Section 3.0 of the SBA.

As part of the South Fork Payette River Subbasin Assessment, the cold water aquatic life beneficial use support status was also determined at fifty-three (53) Beneficial Use Reconnaissance Program monitoring sites encompassing thirty-five (35) streams. Of the 35 streams, it was determined that cold water aquatic life is not fully supported in four of them. These streams may be recommended for §303(d) listing dependant upon results of 2004 BURP data collection. Further data collection efforts may be undertaken, dependent upon results and funding constraints.

Table A summarizes the outcome of the South Fork Payette River Subbasin Assessment.

Table A. Summary of assessment outcomes

Water Body Segment	Pollutant	TMDL(s) Completed	Recommended Changes to §303(d) List	Recommended Schedule Changes
South Fork Payette River WQLS:5186 AU:SW001_05	Sediment	None	De-list sediment	None
Wash Creek – lower WQLS:xxxx AU:SW001_02	Unknown	None	May list pending 2004 BURP data	May prepare informational TMDL
Horn Creek WQLS:xxxx AU:SW001_02	Unknown	None	May list pending 2004 BURP data	May prepare informational TMDL
Chapman Creek WQLS:xxxx AU:SW001_02	Unknown	None	May list pending 2004 BURP data	May prepare informational TMDL
Smokey Creek WQLS:xxxx AU:SW001_02	Unknown	None	May list pending 2004 BURP data	May prepare informational TMDL